

APPROXIMATION OF FUNCTIONS OF TWO VARIABLES ON THE UNIT BICIRCLE

The approximative properties of interpolational Lagrange polynomials, Fourier series of functions of two variables which are defined on the unit bicircle, is stated. For functions from generalized Hölder spaces H_ω the estimations are conducted for norms of C , H_ω , L_p , $p > 1$ spaces [1].

Let γ_{01} and γ_{02} denote the unit circles with centers in the origin; and $\gamma_0 = \gamma_{01} \times \gamma_{02}$ is the unit bicircle. Let $\omega(\delta_1, \delta_2)$ be the some modulus of continuity, and $\Omega_1(\delta)$, $\Omega_2(\delta)$ are the simple moduli of continuity, which respect to that one; $\omega_{1,1}$ denotes the mixed modulus of continuity of the second order [2]. Let H_ω denotes the space of continuous functions $x(t, \tau)$ at γ_0 , which satisfy the conditions:

$$H(x; \omega) = \sup_{\delta_1^2 + \delta_2^2 \neq 0} \frac{\omega(\delta_1, \delta_2; x)}{\omega(\delta_1, \delta_2)} \leq c_1, \quad H^{l\tau}(x; \omega) = \sup_{\delta_1^2 + \delta_2^2 \neq 0} \frac{\omega_{1,1}(\delta_1, \delta_2; x)}{\Omega_1(\delta_1)\Omega_2(\delta_2)} \leq c_2.$$

In the space H_ω the norm is defined as:

$$\|x(t, \tau)\|_{H_\omega} = \|x(t, \tau)\|_C + H(x; \omega) + H^{l\tau}(x; \omega).$$

$$\text{Let } (L_{mn}x)(t, \tau) = \sum_{p=0}^{2m} \sum_{q=0}^{2n} x(t_p, \tau_q) l_p(t) l_q(\tau) = \sum_{k=-m}^m \sum_{l=-n}^n \lambda_{kl} t^k \tau^l,$$

$$l_p(t) = \prod_{\substack{k=0 \\ k \neq p}}^{2m} \frac{t - t_k}{t_p - t_k} \left(\frac{t_p}{t} \right)^m = \sum_{k=-m}^m \lambda_{1,k}^{(p)} t^k, \quad l_q(\tau) = \prod_{\substack{j=0 \\ j \neq q}}^{2n} \frac{\tau - \tau_j}{\tau_q - \tau_j} \left(\frac{\tau_q}{\tau} \right)^n = \sum_{j=-n}^n \lambda_{2,j}^{(q)} \tau^j,$$

denotes interpolational Lagrange polynomial of the function $x(t, \tau)$ with respect to the system of equidistant points at γ_0 :

$$t_p = \exp\left\{\frac{2\pi i}{2m+1}(p-m)\right\}, \quad i^2 = -1, \quad p = 0, \dots, 2m;$$

$$\tau_q = \exp\left\{\frac{2\pi i}{2n+1}(q-n)\right\}, \quad i^2 = -1, \quad q = 0, \dots, 2n.$$

Let operator Φ_{mn} assigns to any function $x(t, \tau) \in H_\omega$ the mn -th partial sum of its Fourier series, that is,

$$(\Phi_{mn}x)(t, \tau) = \sum_{k=-m}^m \sum_{l=-n}^n x_{kl} t^k \tau^l; \quad x_{kl} = \frac{1}{4\pi^2} \int_{\gamma_{01}} \int_{\gamma_{02}} \frac{x(t, \tau)}{t^{k+1} \tau^{l+1}} dt d\tau, \quad k, l = 0, \pm 1, \pm 2, \dots$$

Theorem 1. For any function $x(t, \tau) \in H_\omega$ and for arbitrary natural m, n the inequality is fulfilled:

$$\|x - L_{mn}x\|_C \leq c l n m l n n H(x; \omega) \omega\left(\frac{1}{m}, \frac{1}{n}\right);$$

$$\|x - \Phi_{mn}x\|_C \leq (1 + c l n (2m+1) n (2n+1)) H(x; \omega) \omega\left(\frac{1}{m}, \frac{1}{n}\right).$$

Theorem 2. Let $x(t, \tau) \in H_{\omega^{(1)}}$; $\omega^{(2)}(\delta_1, \delta_2)$ is such that $H_{\omega^{(1)}} \subset H_{\omega^{(2)}}$ and $\Omega_1^{(1)}(\delta)/\Omega_1^{(2)}(\delta)$, $\Omega_2^{(1)}(\delta)/\Omega_2^{(2)}(\delta)$ are increasing functions [3]. Then:

$$\|x - L_{mn}x\|_{H_{\omega^{(2)}}} \leq c l n m l n n \frac{\omega^{(1)}\left(\frac{1}{m}, \frac{1}{n}\right)}{\Omega_1^{(2)}\left(\frac{1}{m}\right) \Omega_2^{(2)}\left(\frac{1}{n}\right)} \|x\|_{H_{\omega^{(1)}}};$$

$$\|x - \Phi_{mn}x\|_{H_{\omega^{(2)}}} \leq c l n m l n n \frac{\omega^{(1)}\left(\frac{1}{m}, \frac{1}{n}\right)}{\Omega_1^{(2)}\left(\frac{1}{m}\right) \Omega_2^{(2)}\left(\frac{1}{n}\right)} \|x\|_{H_{\omega^{(1)}}}.$$

Theorem 3. Let $x(t, \tau) \in H_\omega$. Then the estimation is fulfilled ($p > 1$):

$$\|x - L_{mn}x\|_{L_p} \leq (1 + c(p)) H(x; \omega) \omega\left(\frac{1}{m}, \frac{1}{n}\right),$$

where $\|L_{mn}\|_{C \rightarrow L_p} \leq c(p)$.

REFERENCES

1. Snizhko, N. Approximation of functions of two variables on the unit bicircle in generalized Hölder spaces [Текст] / N. Snizhko // Abstracts of VIII International Scientific and Practical Conference. Prague, Czech Republic, 2020. – P. 577–582

2. Тиман, А. Ф. Теория приближения функций действительного переменного [Текст] / А. Ф. Тиман. – М. : ГИФМЛ, 1965. – 624 с.

3. Сніжко, Н. В. Класифікація узагальнених просторів Гельдера функцій двох змінних [Текст] / Н. В. Сніжко // Вісник Київського університету. Сер. фіз.-мат. науки. – 1999. – Вип. 3. – С. 124–128.